## **EmONC** signal function checklists

Lists of EmONC signal functions by potential EmONC level:

Basic EmONC Signal Functions	Performed	Not performed
OBSTETRIC		
Administer medications to treat postpartum hemorrhage		
Administer parenteral antibiotics (woman)		
Administer magnesium sulfate		
Remove retained products of conception		
Perform manual removal of placenta		
Perform assisted vaginal birth		
Provide IV fluid replacement therapy (woman)		
NEONATAL		
Perform newborn resuscitation with bag and mask		
Initiate and support early and exclusive breastfeeding		
Administer parenteral antibiotics (newborn)		
Practice immediate kangaroo mother care for preterm and low birthweight infants		
Administer oxygen therapy with pulse oximetry for stabilization and transportation (newborn)		
REFERRAL		
Arrange ambulance, with trained and equipped provider, to a facility that can provide definitive care (woman and/or newborn)		

Comprehensive EmONC Signal Functions	Performed	Not performed
OBSTETRIC		
Administer medications to treat postpartum hemorrhage		
Administer parenteral antibiotics (woman)		
Administer magnesium sulfate		
Remove retained products of conception		
Perform manual removal of placenta		
Perform assisted vaginal birth		
Provide IV fluid replacement therapy (woman)		
Administer antenatal corticosteroids		
Perform cesarean section		
Perform blood transfusion (woman)		
NEONATAL		
Perform newborn resuscitation with bag and mask		
Initiate and support early and exclusive breastfeeding		
Administer parenteral antibiotics (newborn)		
Practice immediate kangaroo mother care for preterm and low birthweight infants		
Provide thermal care with radiant warmer or incubator (newborn)		
Administer oxygen therapy with pulse oximetry (newborn)		
Provide CPAP (newborn)		
Provide phototherapy (newborn)		
Perform blood transfusion (newborn)		
Enable assisted feeding with expressed breast milk with cup and spoon and tube feeding		
Provide IV fluids (newborn)		

Intensive EmONC Signal Functions	Performed	Not performed
OBSTETRIC		
Administer medications to treat postpartum hemorrhage		
Administer parenteral antibiotics (woman)		
Administer magnesium sulfate		
Remove retained products of conception		
Perform manual removal of placenta		
Perform assisted vaginal birth		
Provide IV fluid replacement therapy (woman)		
Administer antenatal corticosteroids		
Perform cesarean section		
Perform blood transfusion (woman)		
Provide intensive level organ support (woman)		
NEONATAL		
Perform newborn resuscitation with bag and mask		
Initiate and support early and exclusive breastfeeding		
Administer parenteral antibiotics (newborn)		
Practice immediate kangaroo mother care for preterm and low birthweight infants		
Provide thermal care with radiant warmer or incubator (newborn)		
Administer oxygen therapy with pulse oximetry (newborn)		
Provide CPAP (newborn)		
Provide phototherapy (newborn)		
Perform blood transfusion (newborn)		
Enable assisted feeding with expressed breast milk with cup and spoon and tube feeding		
Provide IV fluids (newborn)		
Perform mechanical ventilation (newborn)		
Perform screening and treatment for retinopathy of prematurity (newborn)		